

Additions and corrections to the list of Norwegian Dixidae (Diptera)

ØYVIND HÅLAND

Håland, Ø. 2013. Additions and corrections to the checklist of Norwegian Dixidae (Diptera). *Norwegian Journal of Entomology* 60, 137–139.

Dixella autumnalis (Meigen, 1838) is recorded for the first time from Norway, from the island of Hvasser in Vestfold, southeastern Norway. A picture of the male hypopygium is given. *Dixa nubilipennis* Curtis, 1832 and *Dixa submaculata* Edwards, 1920 are deleted from the list as the records were based on misidentifications. *Dixa maculata* Meigen, 1818 is recorded for the second time from Norway, from southern Hedmark. An updated list of the Norwegian Dixidae is presented.

Key words: Diptera, Dixidae, *Dixella autumnalis*, Norway, distribution, check list.

Øyvind Håland, Kvalvegen 2, NO-2385 Brumunddal, Norway. E-mail: oyvhalan@broadpark.no

Introduction

Three larvae of a dixid species, brought to my attention by Kjell Magne Olsen, was with some hesitation identified to *Dixella autumnalis* (Meigen, 1838). However, the setae on the ventral side of the caudal filament were not arranged according to the table in Disney (1999). On 1 July 2012 the locality was therefore revisited and more larvae were collected for rearing. One of these larvae did pupate and a male hatched on 12 July 2012.

The specimen was mounted on a slide in Euparal and could then be identified with certainty to *D. autumnalis*; the hypopygium is shown in Figure 1. The locality is a small, shallow pond, situated about 2m a.s.l. in a Nature Reserve at the southern tip of Tjønneberget on the island of Hvasser, (VE, Tjøme: Hvasser, Tjønneberget, N 59°06413, E 10°43765, EIS 19). The area is used as a grazing area for sheep. The water in the pond is probably slightly saline, being very close to the sea. It was found together with larvae of *Dixella amphibia* (De Geer, 1776) and *Dixella aestivalis* (Meigen, 1818).

D. autumnalis has previously been recorded from The Åland Islands (Finland) (Salmela 2006, Autio & Salmela 2010) and is reported to be common in the lowlands on the British Islands (Disney 1999) and in The Netherlands (Krebs 1982). It seems to tolerate some salinity, viz. 3.4‰ (Krebs 1982). According to Wagner (2013) the European distribution covers: Belgium, Corsica, Denmark, Finland, France, Great Britain, Germany, Ireland, Italy, Sardinia, Spain, Switzerland, and The Netherlands.

Dixa nubilipennis Curtis, 1832, was recorded from a brook near Byflaten north of Brumunddal in Ringsaker municipality in southern Hedmark by Olsen (2008). Later another record from a nearby locality was given by Håland (2011); this specimen was sent to Dr. J.K. Moulton at the University of Tennessee (USA) for genetic analysis. The first specimen was recently re-examined and identified to *Dixa maculata* (Meigen, 1818), a species that previously has been recorded only once from Norway, from Tøyen, Oslo in 1841 (Peus 1936). For the time being it is impossible to separate the larvae of *Dixa nubilipennis* and *D. maculata* and the male hypopygium of *D. maculata* has to be



FIGURE 1. Hypopygium of the male of *Dixella autumnalis* (Meigen, 1838).

cleared in KOH to reveal the sclerotized internal hooks to allow identification.

Dixa submaculata Edwards, 1920, was recorded by Håland (1996) from Sandnessjøen in Nordland, based on a few larvae. At closer examination they apparently were misidentified and the species must be deleted from the list of Norwegian Dixidae. However, *D. submaculata* may well occur in Norway as it is recorded from Sweden (Tjeder 1954) and Finland (Salmela 2006).

Check list

A revised list of Norwegian Dixidae is given below, consisting of 17 species.

- *Dixa dilatata* Strobl, 1900
- *Dixa maculata* Meigen, 1818
- *Dixa nebulosa* Meigen, 1830
- *Dixa puberula* Loew, 1849
- *Dixella aestivalis* (Meigen, 1818)
- *Dixella amphibia* (De Geer, 1776)
- *Dixella attica* (Pandazis, 1933)
- *Dixella autumnalis* (Meigen, 1838)
- *Dixella borealis* (Martini, 1928)
- *Dixella dyari* (Garrett, 1924)
- *Dixella filicornis* (Edwards, 1926)
- *Dixella hyperborea* (Bergroth, 1889)

- *Dixella martini* (Peus, 1934)
- *Dixella naevia* (Peus, 1934)
- *Dixella nigra* (Staeger, 1840)
- *Dixella obscura* (Loew, 1849)
- *Dixella serotina* (Meigen, 1818)

Discussion

A few more species are known to occur in our neighboring countries (see Wagner 2013), and some of them might well be found in Norway too. Further, J.K. Moulton has recently found several cryptic species among the North American Dixidae species using DNA methods (see Midgepeet 2011). This may well be the case for some European species too, so the number of species known to occur in Norway will probably rise in the years to come.

Acknowledgements. Many thanks are due to Mr. Kjell Magne Olsen for sending me larvae of *Dixella autumnalis* and encouraging me to visit the locality on Hvasser. Many thanks also to førsteamanuensis Trond Andersen at the Bergen Museum for correcting the manuscript.

References

- Autio, O. & Salmela, J. 2010. The semi-aquatic fly fauna (Diptera) of wetlands of the Åland Islands. *Memoranda Societas pro Fauna et Flora Fennica* 86, 43–53.
- Disney, R.H.L. 1999. British Dixidae (meniscus midges) and Thaumaleidae (trickle midges): keys with ecological notes. *Scientific Publications of the Freshwater Biological Association* 56, 1–129.
- Håland, Ø. 1997. *Dixidae*. Pp. 199–201 in Aagaard, K. & Dolmen, D. (Eds.), *Limnofauna norvegica*. Tapir Forlag, Trondheim.
- Håland, Ø. 2011. First Norwegian record of *Dixella naevia* (Peus, 1934) (Dixidae), and other interesting records of nematocerous Diptera. *Norwegian Journal of Entomology* 58, 9–14.
- Krebs, B.P.M. 1982. Note on the distribution of *Dixella autumnalis* (Meigen) in the south-west Netherlands (Diptera: Dixidae). *Entomologische Berichten* 42, 62–64.
- Midgepeet, 2011. A collaborative effort to increase

- taxonomic expertise in understudied families of nematoceros Diptera. Available from: <http://www.midgepeet.utk.edu/dixidae.htm> (Accessed: 9 May 2013).
- Nicholson, P.A. 1978. Some observations on *Dixella autumnalis* Meigen (Dipt., Dixidae). *Entomologist's monthly Magazine* 113, 62.
- Peus, F. 1936. Zur Kenntnis der Dixiden-Fauna Nordeuropas. *Norsk Entomologisk Tidsskrift* 4, 117–127.
- Salmela, J. 2006: Suomen vaaksiaiset, kummitussääsket, perhossääsket, sinkilähyttiset ja norosääsket (Diptera, Nematocera) – ekologia, levinneisyys ja uhanalaisuus. Alustava raportti. [Ecology, distribution and redlist status of craneflies, phantom midges, mothflies, meniscus midges and trickle midges in Finland (Diptera, Nematocera). Preliminary report]. Privately published, 75 pp. In Finnish with English summary.
- Tjeder, B. 1954. Östskånska insekter 2. *Opuscula Entomologica* 19, 205–209.
- Wagner, R. 2013. *Fauna Europaea: Dixidae*. In Beuk, P. & Pape, T. (Eds), *Fauna Europaea: Diptera, Nematocera*. *Fauna Europaea* version 2.6. Available from: <http://www.faunaeur.org> (Accessed: 9 May 2013).

Received: 23 November 2012

Accepted: 22 June 2013