

Taxonomy of some African species hitherto placed in *Stenentoma* Diakonoff, 1969 and in *Eucosmocydia* Diakonoff, 1988 (Lepidoptera, Tortricidae)

LEIF AARVIK & DAVID J.L. AGASSIZ

Aarvik, L. & Agassiz, D.J.L. 2014. Taxonomy of some African species hitherto placed in *Stenentoma* Diakonoff, 1969 and in *Eucosmocydia* Diakonoff, 1988 (Lepidoptera, Tortricidae). *Norwegian Journal of Entomology* 61, 27–36.

Stenentoma Diakonoff, 1969 is synonymized with *Camptrodoxa* Meyrick, 1925. The following species hitherto placed in *Stenentoma* are transferred to *Camptrodoxa*; *C. bisecta* (Meyrick, 1918) **comb. nov.**, *C. plectocosma* (Meyrick, 1921) **comb. nov.**, *C. chrysolampra* (Diakonoff, 1969) **comb. nov.**, *C. onychosema* (Diakonoff, 1969) **comb. nov.**, and *C. sorindeiae* (Razowski & Brown, 2012) **comb. nov.** The new genus *Afroicelita* **gen. nov.** is established for *Stenentoma pholicosta* Razowski & Wojtusiak, 2012. *Camptrodoxa inclyta* Meyrick, 1925 is a junior synonym of *Laspeyresia plectocosma* Meyrick, 1921 **syn. nov.** *Eucosmocydia monitrix* (Meyrick, 1909) is transferred to the genus *Namasia* Diakonoff, 1983, and the valid name for the species is established as *Namasia monitrix* (Meyrick, 1909) **comb. nov.** *Neonamasia* **gen. nov.** is proposed for *Neonamasia cryptica* **sp. n.**

Key words: Lepidoptera, Tortricidae, *Stenentoma*, *Camptrodoxa*, *Namasia*, new genera, new species, Africa.

Leif Aarvik, Natural History Museum, University of Oslo, P.O. Box 1172 Blindern, NO-0318 Oslo, Norway. E-mail: leif.aarvik@nhm.uio.no

David J.L. Agassiz, Terrestrial Invertebrates Division, Natural History Museum, London SW7 5BD, United Kingdom. E-mail: d.agassiz@nhm.ac.uk

Introduction

Investigation of African Tortricidae species currently assigned to *Stenentoma* Diakonoff, 1969 and *Eucosmocydia* Diakonoff, 1988 has led to new generic combinations and synonymy. The little used generic names *Camptrodoxa* Meyrick, 1925 and *Namasia* Diakonoff, 1983 are involved. They were both described from single specimens. The moths are quite small, wingspan 9–12mm, and are recognizable by having a notch at the forewing termen just above middle where vein M3 and the curved vein CuA1 reach the edge. Initially members of this group were thought to

fall into two genera, but an additional undescribed species turned out to represent a new genus. The forewing character links the three genera with the Afrotropical *Thylacogaster* Diakonoff, 1988 and the Neotropical *Eriosocia* Razowski & Brown, 2008. Razowski & Wojtusiak (2012) described *Stenentoma pholicosta* (Razowski & Wojtusiak, 2012) from females from Nigeria. After examination of additional material, including males, from several African countries, it was found that *S. pholicosta* is not closely related to other species placed in *Stenentoma*. A new genus is proposed for the species, and it is placed in the tribe Eucosmini.