

Sawflies (Hymenoptera, Symphyta) newly recorded from Norway

ERIK HEIBO, OLE JØRGEN LØNNVE, TROND ELLING BARSTAD, STEPHAN MARTIN BLANK,
ANDREW LISTON, MARKO PROUS & ANDREAS TAEGER

Heibo, E., Lønnve, O.J., Barstad, T.E., Blank, S.M., Liston, A., Prous, M., & Taeger, A. 2014. Sawflies (Hymenoptera, Symphyta) newly recorded from Norway. *Norwegian Journal of Entomology* 61, 15–26.

22 species are newly recorded in Norway: *Cladardis hartigi* Liston, 1995; *Cladius rufipes* Serville, 1823; *Dineura stilata* (Klug, 1816); *Dolerus (Poodolerus) varispinus* Hartig, 1837; *Empria basalis* Lindqvist, 1968; *Empria camtschatica* Forsius, 1928; *Empria excisa* (Thomson, 1871); *Empria minuta* Lindqvist, 1968; *Eutomostethus punctatus* (Konow, 1887); *Fenusella wuestneii* (Konow, 1895); *Gilpinia virens* (Klug, 1812); *Phylloecus linearis* (Schrank, 1781); *Heterarthrus nemoratus* (Fallén, 1808); *Hoplocampa crataegi* (Klug, 1816); *Hoplocampa flava* (Linné, 1760); *Metallus albipes* (Cameron, 1875); *Nematinus steini* Blank, 1998; *Nematus cadderensis* Cameron, 1875; *Pamphilius marginatus* (Serville, 1823); *Pseudodineura enslini* (Hering, 1923); *Pseudodineura fuscula* (Klug, 1816); *Tenthredo semicolon* Mol, 2013. Comments are made on the European distribution. Hostplants of the presented sawflies are remarked upon in the light of the distribution of the plants in Norway.

Key words: Cephidae, Pamphiliidae, Diprionidae, Tenthredinidae, new records.

* Erik Heibo, Ento Consulting, Rypeveien 34A, NO-3420 Lierskogen, Norway.
E-mail: eheibo@yahoo.no

Ole J. Lønnve, BioFokus, Gaustadaléen 21, NO-0349 Oslo, Norway. E-mail: ole@biofokus.no

Trond Elling Barstad, Department of Natural Sciences, Tromsø University Museum, NO-9037, Tromsø, Norway. E-mail: trond.elling.barstad@uit.no

Stephan M. Blank, Andrew Liston, Marko Prous & Andreas Taeger, Senckenberg Deutsches Entomologisches Institut, Eberswalder Str. 90, DE-15374 Müncheberg, Germany.
E-mails: stephan.blank@senckenberg.de; andrew.liston@senckenberg.de;
marko.prous@senckenberg.de; andreas.taeger@senckenberg.de

* Corresponding author

Introduction

Siebke (1880), Strand (1898) and Kiær (1898) published important works on the Symphyta of Norway, with smaller additions by Strand (1900, 1901, 1903). Since then, sawfly research in Norway has not been continuously performed and until the last decade, few articles containing new records were published (e.g. Heibo & Lønnve

2005, Nuorteva *et al.* 2005, Lønnve 2006, Lønnve 2007, Heibo *et al.* 2008, and Lønnve 2009).

All sawfly larvae are herbivorous (Wagner & Raffa 1993, Liston 1995, Viitassari 2002), except for those of the Orussidae, which are ecto- and endoparasitic on Buprestidae, Cerambycidae and Siricidae (Kraus 1998). Among the plant feeding sawflies both monophagy and oligophagy are common, while polyphagy is probably not that